


EVOLUTION OF THE H7N9 VIRUS IN CHINA, 2013







Tetrapodophis amplectus



Tetrapodophis amplectus


ตัวอย่างฟอสซิลที่มีชีวิต (Living fossil)



ตัวอย่างฟอสซิลที่มีชีวิต (Living fossil)





ตัวอย่างฟอสซิลที่มีชีวิต (Living fossil)



หลักฐานของการวิวัฒนาการ


• Biogeography evidence







หลักฐานของการวิวัฒนาการ

• Homologous structure VS Analogous structure




Homologous structure








หลักฐานของการวิวัฒนาการ


Molecular record















Founder effect





The effects of different types of selection on a population










Figure 8-26 What Is Life? A Guide to Biology © 2010 W.H. Freeman and Company




(a) Normal red blood cell




Normal hemoglobin




(b) Sickled red blood cell



Sickle-cell hemoglobin








Figure 8-27
What Is Life? A Guide to Biology
© 2010 W.H. Freeman and Company





Figure 8-28
What Is Life? A Guide to Biology
© 2010 W.H. Freeman and Company





Horse

Donkey



Mule (hybrid)



Greater Canada goose (Branta canadensis)



Greylag goose (Anser anser)



Abert's squirrel

Kaibab squirrel



PREZYGOTIC BARRIERS

Habitat isolation: populations live in different habitats and do not meet

Behavioral isolation: little or no sexual attraction between males and females

Temporal isolation: mating or flowering occurs at different seasons or times of day

Mating

0

Mechanical isolation: structural differences in genitalia or flowers prevent copulation or pollen transfer

Gametic isolation: female and male gametes fail to attract each other or are inviable

Fertilization


POSTZYGOTIC BARRIERS


Reduced hybrid viability: hybrid zygotes fail to develop or fail to reach sexual maturity

Reduced hybrid fertility: hybrids fail to produce functional gametes

Hybrid breakdown: offspring of hybrids have reduced viability or fertility









PREZYGOTIC BARRIERS

Habitat isolation: populations live in different habitats and do not meet

Behavioral isolation: little or no sexual attraction between males and females

Temporal isolation: mating or flowering occurs at different seasons or times of day

Mating

0

Mechanical isolation: structural differences in genitalia or flowers prevent copulation or pollen transfer

Gametic isolation: female and male gametes fail to attract each other or are inviable

Fertilization

POSTZYGOTIC BARRIERS


Reduced hybrid viability: hybrid zygotes fail to develop or fail to reach sexual maturity

Reduced hybrid fertility: hybrids fail to produce functional gametes

Hybrid breakdown: offspring of hybrids have reduced viability or fertility









Mate late winter

Viable, fertile offspring



PREZYGOTIC BARRIERS

Habitat isolation: populations live in different habitats and do not meet

Behavioral isolation: little or no sexual attraction between males and females

Temporal isolation: mating or flowering occurs at different seasons or times of day

Mating

0

Mechanical isolation: structural differences in genitalia or flowers prevent copulation or pollen transfer

Gametic isolation: female and male gametes fail to attract each other or are inviable


Fertilization

POSTZYGOTIC BARRIERS

Reduced hybrid viability: hybrid zygotes fail to develop or fail to reach sexual maturity

Reduced hybrid fertility: hybrids fail to produce functional gametes







PREZYGOTIC BARRIERS

Habitat isolation: populations live in different habitats and do not meet

Behavioral isolation: little or no sexual attraction between males and females

Temporal isolation: mating or flowering occurs at different seasons or times of day

Mating

Mechanical isolation: structural differences in genitalia or flowers prevent copulation or pollen transfer

Gametic isolation: female and male gametes fail to attract each other or are inviable


Fertilization


POSTZYGOTIC BARRIERS

Reduced hybrid viability: hybrid zygotes fail to develop or fail to reach sexual maturity

Reduced hybrid fertility: hybrids fail to produce functional gametes









Habitat isolation: populations live in different habitats and do not meet

Behavioral isolation: little or no sexual attraction between males and females

Temporal isolation: mating or flowering occurs at different seasons or times of day

Mating

0

Mechanical isolation: structural differences in genitalia or flowers prevent copulation or pollen transfer

Gametic isolation: female and male gametes fail to attract each other or are inviable

Fertilization

POSTZYGOTIC BARRIERS

Reduced hybrid viability: hybrid zygotes fail to develop or fail to reach sexual maturity

Reduced hybrid fertility: hybrids fail to produce functional gametes











PREZYGOTIC BARRIERS

Habitat isolation: populations live in different habitats and do not meet

Behavioral isolation: little or no sexual attraction between males and females

Temporal isolation: mating or flowering occurs at different seasons or times of day

Mating

0

Mechanical isolation: structural differences in genitalia or flowers prevent copulation or pollen transfer

Gametic isolation: female and male gametes fail to attract each other or are inviable


Fertilization


POSTZYGOTIC BARRIERS

Reduced hybrid viability: hybrid zygotes fail to develop or fail to reach sexual maturity

Reduced hybrid fertility: hybrids fail to produce functional gametes







การปรับตัว

• Deceptive coloration




การปรับตัว

• Warning (aposematic) coloration






วิวัฒนาการของพืช







(b) Carpel evolution

วิวัฒนาการของสัตว์

